

JEEM

Journal of Embryology and Experimental Morphology

Volume 77 October 1983

| | |
|---|---------|
| Lim, S.-S., Woodroffe, M. N. and Lemanski, L. F. An analysis of contractile proteins in developing chick heart by SDS polyacrylamide gel electrophoresis and electron microscopy | 1-14 |
| Ubbels, G. A., Hara, K., Koster, C. H. and Kirschner, M. W. Evidence for a functional role of the cytoskeleton in determination of the dorsoventral axis in <i>Xenopus laevis</i> eggs | 15-37 |
| Schmidt, J. T. Regeneration of the retinotectal projection following compression onto a half tectum in goldfish | 39-51 |
| Cooke, J. and Gaze, R. M. The positional coding system in the early eye rudiment of <i>Xenopus laevis</i> , and its modification after grafting operations | 53-71 |
| Stern, C. D. and MacKenzie, D. O. Sodium transport and the control of epiblast polarity in the early chick embryo | 73-98 |
| Beresford, B. Brachial muscles in the chick embryo: the fate of individual somites | 99-116 |
| Ashwell, K. W. and Watson, C. R. R. The development of facial motoneurons in the mouse - neuronal death and the innervation of the facial muscles | 117-141 |
| Azar, Y. and Eyal-Giladi, H. The retention of primary hypoblastic cells underneath developing primitive streak allows for their prolonged inductive influence | 143-151 |
| Fucci, L., Cirotto, C., Tomei, L. and Geraci, G. Synthesis of globin chains in the erythropoietic sites of the early chick embryo | 153-165 |
| Graziosi, G., De Cristini, F., Di Marcotullio, A., Marzari, R., Micali, F. and Savoini, A. Morphological and molecular modifications induced by heat shock in <i>Drosophila melanogaster</i> embryos | 167-182 |
| Gualandris, L., Rougé, P. and Duprat, A. M. Membrane changes in neural target cells studied with fluorescent lectin probes | 183-200 |
| De Pomerai, D. I., Kotecha, B., Flor-Henry, M., Fullick, C., Young, A. and Gali, M. A. H. Expression of differentiation markers by chick embryo neuroretinal cells <i>in vivo</i> and in culture | 201-220 |
| Reynolds, S., Holder, N. and Fernandes, M. The form and structure of supernumerary hindlimbs formed following skin grafting and nerve deviation in the newt <i>Triturus cristatus</i> | 221-241 |
| Sawai, T. Cytoplasmic and cortical factors participating in cleavage furrow formation in eggs of three amphibian genera; <i>Ambystoma</i> , <i>Xenopus</i> and <i>Cynops</i> | 243-254 |
| Schrag, J. A. and Cameron, J. A. Regeneration of adult newt skeletal muscle tissue <i>in vitro</i> | 255-271 |
| Maden, M. The effect of vitamin A on the regenerating axolotl limb | 273-295 |
| MacQueen, H. A. and Johnson, M. H. The fifth cell cycle of the mouse embryo is longer for smaller cells than for larger cells | 297-308 |
| Vasse, J. Transplantation of turtle embryonic thymus into quail embryo: colonization by quail cells | 309-322 |
| Index of Authors and Titles | 323-324 |
| Subject Index | 325-327 |

ISSN: 0022-0752

The Company of Biologists Limited
Department of Zoology, University of Cambridge, Downing Street,
Cambridge CB2 3EJ

© The Company of Biologists Limited 1983

Filmset by Filmtyp Services Limited and Printed in England by Pindar Print Limited,
Scarborough, North Yorkshire.